



## Synthesis and Theoretical Study of Zinc(II) and Nickel(II)-Complexes of 5-Methoxyisatin 3-[N-(4-chlorophenyl)thiosemicarbazone]

Fatma Kandemirli<sup>1</sup>, Murat Saracoğlu<sup>2</sup>, Gamze Bulut<sup>3</sup>, Eno Ebenso<sup>4</sup>,  
Taner Arslan<sup>5</sup> & Asgar Kayan<sup>3</sup>

<sup>1</sup>Department of Chemistry, Niğde University, 51240, Niğde, Turkey

<sup>2</sup>Faculty of Education, Erciyes University, 38039, Kayseri, Turkey

<sup>3</sup>Department of Chemistry, Kocaeli University, 41380, Izmit, Turkey

<sup>4</sup>Department of Chemistry, North West University (Mafikeng Campus), Private Bag X2046, Mmabatho 2735, South Africa

<sup>5</sup>Department of Chemistry, Osmangazi University, 26480, Eskişehir, Turkey  
Email: fkandemirli@yahoo.com

**Abstract.** Zinc(II) and nickel(II)-complexes of 5-methoxyisatin 3-[N-(4-chlorophenyl) thiosemicarbazone] (H<sub>2</sub>MICP) were synthesized and characterized by infrared, ultraviolet and <sup>1</sup>H-NMR spectroscopies as well as elemental analysis. Model of H<sub>2</sub>MICP and its zinc(II) and nickel(II)-complexes were optimized with B3LYP method using 6-31G(d,p), 6-311G(d,p), 6-311++G(d,p), 6-311++G(2d,2p) basis sets. The calculated <sup>1</sup>H-NMR, UV and IR spectra data were compared with experimental results. In addition to the Natural Bond Orbital (NBO) analysis of H<sub>2</sub>MICP and its Zinc(II) and Nickel(II) complexes, Fukui functions of H<sub>2</sub>MICP were also reported.

**Keywords:** DFT; isatin thiosemicarbazones; Zn (II) and Ni (II)-complexes.

### 1 Introduction

Isatin 3-thiosemicarbazone derivatives and their metal complexes have a broad range of biological activities namely antimicrobial [1], antiviral [2,3], antitumor, anti-inflammatory and antibacterial activities [4]. The investigation of the structure-activity relationships of isatin β-thiosemicarbazones for antiviral chemotherapeutic activity revealed that the substitution at the C=O position of the side-chain by another atom or group should result in loss of activity [2]. Recently, structure-activity relationship of 5-fluoro-1*H*-indole-2,3-dione-3-thiosemicarbazones and 5-fluoro-1-morpholino/piperidinomethyl-1*H*-indole-2,3-dione-3-thiosemicarbazones were evaluated for *in vitro* antituberculosis activity against *Mycobacterium tuberculosis* H37Rv and studied using ETM-ANN method [5,6]. Antimicrobial activity of Schiff and Mannich bases derived from isatin derivatives has been reported. *N*-[4-(4'-chlorophenyl) thiazol-2-yl] thiosemicarbazide and 1-[*N,N*-dimethylaminomethyl]-5-bromo isatin-3'-{1'-[4''-(*p*-chlorophenyl)thiazol-2''-

Received September 8<sup>th</sup>, 2011, Revised April 30<sup>th</sup>, 2011, 2<sup>nd</sup> Revision May 6<sup>th</sup>, 2011, 3<sup>rd</sup> Revision October 6<sup>th</sup>, 2011, Accepted for publication November 4<sup>th</sup>, 2011.

Copyright © 2012 Published by LPPM ITB, ISSN: 1978-3043, DOI: 10.5614/itbj.sci.2012.44.1.4

yl] thio semicarbazone showed the most favorable antimicrobial activity [1]. 1-(1-((Substituted)methyl)-5-methyl-2-oxoindolin-3-ylidene)-4-(substituted pyridine-2-yl)thiosemicarbazide has been reported to show significant anti-inflammatory and analgesic activity [1].

There are a number of studies on the synthesis of isatin  $\beta$ -thiosemicarbazone derivatives. We have reported synthesis, characterization and mechanistic of 5-methoxyisatin 3-[N-(4-chlorophenyl)thiosemicarbazone] [7]. In the present study, zinc(II) and nickel(II)-complexes of 5-methoxyisatin 3-[N-(4-chlorophenyl)thiosemicarbazone] have been synthesized. The structures of complexes have been determined by  $^1\text{H-NMR}$ , IR and UV spectra and elemental analysis (C, H, N, S). Moreover, electronics parameters of  $\text{H}_2\text{MICP}$  and its zinc(II) and nickel(II)-complexes have also been calculated using B3LYP with the basis set of 6-31G(d,p), 6-311G(d,p), 6-311++G(d,p), 6-311++G(2d,2p). Natural Bond Orbital (NBO) analysis is also reported.

## 2 Experimental

### 2.1 Material and Methods

5-methoxyisatin was purchased from Sigma Aldrich. Metal salts were purchased from E. Merck and used without further purification. Infrared spectrum of 5-methoxy isatin and its complexes were measured using KBr pellet on Shimadzu FT-IR 8201 spectrometer. The elemental analysis was carried out on CHNS-932 (LECO) and  $^1\text{H-NMR}$  spectra were measured at 400 MHz on a BRUKER DPX-400 spectrometer at the TUBITAK ATAL instrumental analyses laboratory. The electronic spectra of UV–visible zone (200–600 nm) of all the compounds were measured (1 cm quartz cell, 0–2.5 absorbance values range) using Shimadzu UV-1601PC spectrophotometer.

### 2.2 Synthesis

$\text{H}_2\text{MICP}$  ligand molecule was synthesized according to literature procedure [7].

#### **Bis{5-methoxyisatin-3-[N-(4-chlorophenyl)thiosemicarbazonato]}zinc(II) [Zn(HMICP) $_2$ .H $_2$ O]**

[Zn(HMICP) $_2$ ] were synthesized by dissolving 1 mmol (0.361 g)  $\text{H}_2\text{MICP}$  and 0.5 mmol (0.106 g) zinc(II) acetate in warm ethanol. Subsequently, the solution was refluxed for 3 hours. Brown solid was isolated by filtration and washed with 95% ethanol, diethylether and dried in vacuum at room temperature. [(calc. (%): C = 48.95, H = 3.08, N = 14.27, S = 8.16, found. (%):

C = 49.09, H = 3.18, N = 14.24, S = 8.84). <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, ppm): 3.55 (s, CH<sub>3</sub>-methoxy), 6.85-7.66 (aromatic C-H), 10.65 (δ, NH), 10.81 (s, indole-NH)].

**Bis{5-methoxyisatin-3-[N-(4-chlorophenyl)]thiosemicarbazonato}nickel(II) [Ni(HMICP)<sub>2</sub>·H<sub>2</sub>O]**

[Ni(HMICP)<sub>2</sub>] were synthesized by dissolving 1 mmol (0.361 g) H<sub>2</sub>MICP and 0.5 mmol (0.088 g) nickel(II) acetate in warm ethanol. Subsequently, the solution was refluxed for 2 h, and stirred for two days at room temperature. A brown solid was isolated and washed with 95% ethanol and diethylether [(calc. (%): C = 48.26, H = 3.29, N = 14.07, S = 8.05, found. (%): C = 48.13, H = 3.22, N = 13.86, S = 7.31)].

### 3 Theoretical Calculations


All calculations were carried out by employing density functional theory (DFT) with the basis set levels [8-10] of B3LYP/6-31G(d,p), B3LYP/6-311G(d,p), B3LYP/6-311++G(d,p) and B3LYP/6-311++G(2d,2p) for the ligand and B3LYP/6-31G(d,p), B3LYP/6-311G(d,p) for the zinc(II) and nickel(II)-complexes. UV calculation for H<sub>2</sub>MICP ligand was also performed by DFT at the level of BP86/CEP\* using 3/76=1000003000, 3/77=0720007000, 3/78=0810010000, 3/74=406 iop values.

### 4 Results and Discussion


The B3LYP/6-311G(d,p) optimized structures of H<sub>2</sub>MICP and its zinc(II) and nickel(II)-complexes were shown in Figure 1.

The possible tautomeric structures for H<sub>2</sub>MICP were calculated using B3LYP/6-31G(d,p), B3LYP/6-311G(d,p), B3LYP/6-311++G(d,p) and B3LYP/6-311++G(2d,2p) basis sets. There is no negative imaginary frequency observed for possible tautomeric forms, indicating that all structures are true minima. The B3LYP/6-311G(d,p) optimized structures of possible tautomeric forms were shown in Figure 2. As summarized in Table 1, the A form was found to be most stable than the other forms with all calculation methods like the study in [11].

The energy differences between most stable and unstable tautomeric forms were found as 34.78 kcal/mol; 21.21 kcal/mol; 32.50 kcal/mol; 31.16 kcal/mol at B3LYP method with 6-31G(d,p), 6-311G (d,p), 6-311++G(d,p) and 6-311++G(2d,2p) basis sets respectively. The dipole moment of A form is the highest of all calculated basis sets while E<sub>HOMO</sub> of A form is the lowest one.



**Figure 1** The structures of  $H_2MICP$  and its zinc(II) and nickel(II)-complexes optimized with B3LYP/6-311G(d,p).



**Figure 2** The possible tautomer structures of  $H_2MICP$  optimized with B3LYP/6-311G(d,p).

**Table 1** The sum of the zero point and electronic energies (au), polarizability ( $\alpha$ ), dipole moment ( $\mu$ ),  $E_{\text{HOMO}}$ ,  $E_{\text{LUMO}}$ , hardness ( $\eta$ ) and softness ( $\sigma$ ) for the possible tautomer structures of H<sub>2</sub>MICP.

Tautomer structures	6-31G(d,p)	6-311G(d,p)	6-311++G(d,p)	6-311++G(2d,2p)				
Sum of the zero point and electronic energies (au)								
A	-1845.119402	-1845.396917	-1845.414918	-1845.451098				
B	-1845.092834	-1845.370541	-1845.389447	-1845.425976				
C	-1845.087848	-1845.342684	-1845.387149	-1845.423678				
D	-1845.087842	-1845.371603	-1845.391259	-1845.429274				
E	-1845.063968	-1845.342684	-1845.363113	-1845.401429				
F	-1845.067024	-1845.345660	-1845.366705	-1845.404319				
$E_{\text{HOMO}}$ (au)								
A	-0.21650	-0.22428	-0.22803	-0.22688				
B	-0.21305	-0.22142	-0.22558	-0.22430				
C	-0.19856	-0.21499	-0.21096	-0.20994				
D	-0.19856	-0.21901	-0.22329	-0.22234				
E	-0.20648	-0.21499	-0.21967	-0.21868				
F	-0.19656	-0.20484	-0.20951	-0.20844				
$E_{\text{LUMO}}$ (au)								
A	-0.09655	-0.10433	-0.10921	-0.10823				
B	-0.09932	-0.10701	-0.11173	-0.11031				
C	-0.08034	-0.09519	-0.09371	-0.09308				
D	-0.08039	-0.09331	-0.09911	-0.09872				
E	-0.08754	-0.09519	-0.10075	-0.09992				
F	-0.08412	-0.09166	-0.09695	-0.09575				
$\eta$ (au) $\sigma$ (au <sup>-1</sup> )								
A	0.060	8.337	0.060	8.337	0.059	8.416	0.059	8.428
B	0.057	8.793	0.057	8.740	0.057	8.783	0.057	8.773
C	0.059	8.459	0.060	8.347	0.059	8.529	0.058	8.557
D	0.059	8.462	0.063	7.955	0.062	8.053	0.062	8.089
E	0.059	8.408	0.060	8.347	0.059	8.409	0.059	8.420
F	0.056	8.894	0.056	8.835	0.056	8.884	0.056	8.874
$\alpha$ $\mu$ (D)								
A	282.02	9.22	294.54	9.22	316.48	8.98	324.25	8.85
B	285.15	5.64	297.80	5.68	320.43	5.36	327.28	5.19
C	284.49	4.22	289.70	4.75	317.05	4.14	324.34	4.03
D	284.58	4.22	286.02	7.52	307.21	7.27	314.30	7.22
E	279.10	4.64	289.70	4.75	311.12	4.49	317.38	4.39
F	287.68	0.98	298.93	1.01	320.54	1.22	326.97	1.20

Fukui functions of H<sub>2</sub>MICP were calculated using AOMix program [11,12]. Fukui functions give information about the reactive regions for nucleophilic and electrophilic attack.  $f_k^+ = \rho_k(N+1) - \rho_k(N)$  measures the changes as the molecule gains electrons which indicating the reactivity for nucleophilic attack, while  $f_k^- = \rho_k(N) - \rho_k(N-1)$  measures the changes as the molecule losses

electrons which indicating the reactivity for electrophilic attack. Table 2 summarizes the Fukui functions for the HOMO and the LUMO of the molecules.

**Table 2** Neutral and protonated Fukui functions for the HOMO and the LUMO of the H<sub>2</sub>MICP using the different basis sets.

H <sub>2</sub> MICP Atoms	6-31G(d,p)		6-311G(d,p)		6-311++G(d,p)	
	$f_k^-$	$f_k^+$	$f_k^-$	$f_k^+$	$f_k^-$	$f_k^+$
C2	-	-	-	-	1.05	-
C5	3.40	-	3.71	-	2.90	-
C6	11.86	5.68	12.13	5.86	11.06	5.81
N7	8.07	1.64	8.35	1.57	6.61	1.63
S8	15.07	8.37	13.40	8.55	19.78	9.01
O10	6.82	8.12	6.75	7.65	5.95	7.45
C11	14.66	3.10	14.98	3.01	13.25	2.79
O12	9.03	-	9.31	-	7.32	-
C13	9.97	-	10.29	-	8.93	-
C14	1.73	8.22	1.88	9.07	1.80	10.03
C17	6.26	3.43	6.32	3.77	5.66	3.58
C18	-	5.86	-	5.76	1.04	5.13
C19	-	6.31	-	6.61	-	6.86
C20	2.18	13.97	2.15	14.25	2.31	14.09
N21	-	1.83	-	1.75	1.62	1.95
N22	4.38	4.14	4.32	3.96	1.95	4.11
N23	1.76	25.01	1.73	24.47	1.55	24.30

Although the compositions of the HOMO and the LUMO for H<sub>2</sub>MICP depend on the basis set, the changes are not significant. For the HOMO, the contributions are mainly from thiosemicarbazone group (S8: 19.78%, N22: 4.32%, N23: 1.73%) and isatin group (C5: 3.71%, C6: 12.13%, C11: 14.98%, C13: 10.29%, C14: 1.73%, C17: 6.32%, C20: 2.15%, N7: 8.35%, O10: 6.75%, O12: 9.31%).

The NBO program performs the analysis of a many-electron molecular wavefunction in terms of localized electron-pair bonding units. The program carries out the determination of natural atomic orbitals (NAOs), natural hybrid orbitals (NHOs), natural bond orbitals (NBOs), and natural localized molecular orbitals (NLMOs). These parameters are applicable to perform natural population analysis (NPA) [13]. Bond orbital coefficients as well as the hybrids (percents of *s* and *p* character) for H<sub>2</sub>MICP, Zn(II) and Ni(II)-complexes are summarized in Tables 3 and 4.

As shown in Table 3, C6-C13, C5-C11, C17-C19 bonds belonging to the benzene ring of isatin group of H<sub>2</sub>MICP have double bond character and the NBO bonds calculated with B3LYP/6-311++G(2d,2p) are 0.7054 C<sub>6</sub> (sp<sup>1.86</sup>) +

0.7088 C<sub>13</sub> (sp<sup>1.60</sup>) + 0.7249 C<sub>6</sub> (p) + 0.6889 C<sub>13</sub> (p) for C6-C13 bond; 0.6966 C<sub>5</sub> (sp<sup>1.85</sup>) + 0.7175 C<sub>11</sub> (sp<sup>1.53</sup>) + 0.7087 C<sub>5</sub> (p) + 0.7055 C<sub>11</sub> (p) for C5-C11 bond; 0.7196 C<sub>17</sub> (sp<sup>1.62</sup>) + 0.6943 C<sub>19</sub> (sp<sup>1.80</sup>) + 0.7187 C<sub>17</sub> (p) + 0.6953 C<sub>19</sub> (p) for C17-C19 bond. While the calculations which carried out using B3LYP/6-311G(d,p) basis set is 0.7074 C<sub>6</sub> (sp<sup>1.81</sup>) + 0.7068 C<sub>13</sub> (sp<sup>1.60</sup>) + 0.7252 C<sub>6</sub> (p) + 0.6885 C<sub>13</sub> (p), 0.6986 C<sub>5</sub> (sp<sup>1.79</sup>) + 0.7156 C<sub>11</sub> (sp<sup>1.55</sup>) + 0.7090 C<sub>5</sub> (p) + 0.7052 C<sub>11</sub> (p) and 0.7163 C<sub>17</sub> (sp<sup>1.64</sup>) + 0.6978 C<sub>19</sub> (sp<sup>1.75</sup>) + 0.7195 C<sub>17</sub> (p) + 0.6945 C<sub>19</sub> (p). It was shown that there are no significant changes in the calculation results using both the 6-311++G(2d,2p) and 6-311G(d,p) basis sets. Therefore, further calculations were then performed by only the 6-311G(d,p) basis set for the Zn(II) and Ni(II)-complexes. From the NBO analysis, C6-C13, C5-C11, C17-C19 bonds of the benzene ring of isatin group of H<sub>2</sub>MICP have same character for pair bond (Table 4).

NBO bonds of C6-C13, C5-C11, C17-C19 which calculated by B3LYP/6-311G(d,p) basis set are 0.7072 C<sub>6</sub> (sp<sup>1.81</sup>) + 0.7070 C<sub>13</sub> (sp<sup>1.61</sup>) + 0.7220 C<sub>6</sub> (p) + 0.6919 C<sub>13</sub> (p), 0.6893 C<sub>5</sub> (sp<sup>1.79</sup>) + 0.7158 C<sub>11</sub> (sp<sup>1.55</sup>) + 0.7147 C<sub>5</sub> (p) + 0.6994 C<sub>11</sub> (p), 0.7162 C<sub>17</sub> (sp<sup>1.65</sup>) + 0.6979 C<sub>19</sub> (sp<sup>1.78</sup>) + 0.7242 C<sub>17</sub> (p) + 0.6896 C<sub>19</sub> (p), respectively for [Zn(HMICP)<sub>2</sub>] complex and 0.7078 C<sub>6</sub> (sp<sup>1.80</sup>) + 0.7065 C<sub>13</sub> (sp<sup>1.61</sup>) + 0.7210 C<sub>6</sub> (p) + 0.6930 C<sub>13</sub> (p), 0.6988 C<sub>5</sub> (sp<sup>1.78</sup>) + 0.7153 C<sub>11</sub> (sp<sup>1.56</sup>) + 0.7165 C<sub>5</sub> (p) + 0.6976 C<sub>11</sub> (p) and 0.7175 C<sub>17</sub> (sp<sup>1.63</sup>) + 0.6965 C<sub>19</sub> (sp<sup>1.80</sup>) + 0.7310 C<sub>17</sub> (p) + 0.6824 C<sub>19</sub> (p), respectively for [Ni(HMICP)<sub>2</sub>]. There are no significant changes in these bonds for either Zn(II) nor Ni(II)-complexes. C20-N23 NBO of the thiosemicarbazone group of H<sub>2</sub>MICP ligand, and its Zn(II) and Ni(II)-complexes are 0.6326 C<sub>20</sub> (sp<sup>2.00</sup>) + 0.7745 N<sub>23</sub> (sp<sup>1.30</sup>) + 0.6778 C<sub>20</sub> (p) + 0.7353 N<sub>23</sub> (p), 0.6254 C<sub>20</sub> (sp<sup>1.99</sup>) + 0.7803 N<sub>23</sub> (sp<sup>1.34</sup>) + 0.6363 C<sub>20</sub> (p) + 0.7714 N<sub>23</sub> (p) and 0.6370 C<sub>20</sub> (sp<sup>2.02</sup>) + 0.7794 N<sub>23</sub> (sp<sup>1.35</sup>) + 0.6370 C<sub>20</sub> (p) + 0.7708 N<sub>23</sub> (p), respectively. The percentage of C20 atom for H<sub>2</sub>MICP ligand and its Zn(II) and Ni(II)-complexes is 40.02, 39.11 and 39.26. In the formation of σ, s% character of C20 for them is 33.32, 33.40 and 33.13 respectively. s% character of C20 atom for H<sub>2</sub>MICP ligand does not change significantly in the formation of Zn(II) and Ni(II)-complexes. C20% of π NBO bond for H<sub>2</sub>MICP ligand, and its Zn(II) and Ni(II)-complexes is 45.94, 40.49 and 40.58, while those of N23 are 54.06, 59.51 and 59.42, respectively. In the complex formation, bond is polarized (about 59.51% for Zn(II)-complex and about 59.42% for Ni(II)-complex) toward N23. S8-C18 bond having double bond character in the H<sub>2</sub>MICP ligand calculated with B3LYP/6-311++G(2d,2p) method, however it shows single bond character in the Zn(II) and Ni(II)-complexes (NBO bond of S8-C18 is 0.6353 S<sub>8</sub> (sp<sup>4.38</sup>) + 0.7723 C<sub>18</sub> (sp<sup>1.57</sup>) + 0.8278 S<sub>5</sub> (p) + 0.5611 C<sub>18</sub> (p) for H<sub>2</sub>MICP, 0.6994 S<sub>8</sub> (sp<sup>4.80</sup>) + 0.7429 C<sub>18</sub> (sp<sup>2.04</sup>) for Zn(II)-complex, and 0.6638 S<sub>8</sub> (sp<sup>4.96</sup>) + 0.7479 C<sub>18</sub> (sp<sup>2.02</sup>) for Ni(II)-complex)

**Table 3** Calculated NBO analysis data for H<sub>2</sub>MICP calculated with 6-311++G(2d,2p) and 6-311G(d,p) basis set.

Atom numbers	H <sub>2</sub> MICP	
	B3LYP/6-311++G(2d,2p)	Hybrids B3LYP/6-311G(d,p)
C3-C15	0.6991 C <sub>3</sub> (sp <sup>1.85</sup> ) + 0.7151 C <sub>15</sub> (sp <sup>1.83</sup> )	0.7005 C <sub>3</sub> (sp <sup>1.80</sup> ) + 0.7137 C <sub>15</sub> (sp <sup>1.84</sup> )
C15-C16	0.7183 C <sub>15</sub> (sp <sup>1.66</sup> ) + 0.6957 C <sub>16</sub> (sp <sup>1.95</sup> )	0.7163 C <sub>15</sub> (sp <sup>1.64</sup> ) + 0.6978 C <sub>16</sub> (sp <sup>1.75</sup> )
C1-C3	0.7125 C <sub>15</sub> (p) + 0.7017 C <sub>16</sub> (p)	0.7195 C <sub>15</sub> (p) + 0.6945 C <sub>16</sub> (p)
C9-C16	0.7068 C <sub>1</sub> (sp <sup>1.80</sup> ) + 0.7074 C <sub>3</sub> (sp <sup>1.73</sup> )	0.7071 C <sub>1</sub> (sp <sup>1.78</sup> ) + 0.7071 C <sub>3</sub> (sp <sup>1.72</sup> )
C1-C2	0.6978 C <sub>1</sub> (p) + 0.7163 C <sub>3</sub> (p)	0.6975 C <sub>1</sub> (p) + 0.7166 C <sub>3</sub> (p)
C2-C9	0.7080 C <sub>9</sub> (sp <sup>1.79</sup> ) + 0.7062 C <sub>16</sub> (sp <sup>1.80</sup> )	0.7084 C <sub>9</sub> (sp <sup>1.77</sup> ) + 0.7058 C <sub>16</sub> (sp <sup>1.78</sup> )
C2-CI37	0.7048 C <sub>1</sub> (sp <sup>1.84</sup> ) + 0.7094 C <sub>2</sub> (sp <sup>1.60</sup> )	0.7068 C <sub>1</sub> (sp <sup>1.78</sup> ) + 0.7075 C <sub>2</sub> (sp <sup>1.58</sup> )
C6-C13	0.7109 C <sub>2</sub> (sp <sup>1.58</sup> ) + 0.7033 C <sub>9</sub> (sp <sup>1.83</sup> )	0.7109 C <sub>2</sub> (sp <sup>1.56</sup> ) + 0.7033 C <sub>9</sub> (sp <sup>1.78</sup> )
C13-C19	0.6978 C <sub>2</sub> (p) + 0.6806 C <sub>9</sub> (p)	0.7338 C <sub>2</sub> (p) + 0.6794 C <sub>9</sub> (p)
C5-C6	0.6766 C <sub>2</sub> (sp <sup>3.42</sup> ) + 0.7363 Cl <sub>37</sub> (sp <sup>4.68</sup> )	0.6997 C <sub>2</sub> (sp <sup>3.52</sup> ) + 0.7426 Cl <sub>37</sub> (sp <sup>4.40</sup> )
C5-C11	0.7054 C <sub>6</sub> (sp <sup>1.86</sup> ) + 0.7088 C <sub>13</sub> (sp <sup>1.60</sup> )	0.7074 C <sub>6</sub> (sp <sup>1.81</sup> ) + 0.7068 C <sub>13</sub> (sp <sup>1.60</sup> )
C17-C19	0.7249 C <sub>6</sub> (p) + 0.6889 C <sub>13</sub> (p)	0.7252 C <sub>6</sub> (p) + 0.6885 C <sub>13</sub> (p)
C11-C17	0.7100 C <sub>9</sub> (sp <sup>1.73</sup> ) + 0.7042 C <sub>19</sub> (sp <sup>1.88</sup> )	0.7066 C <sub>9</sub> (sp <sup>1.75</sup> ) + 0.7076 C <sub>19</sub> (sp <sup>1.84</sup> )
N7-C11	0.7087 C <sub>5</sub> (sp <sup>1.78</sup> ) + 0.7055 C <sub>6</sub> (sp <sup>1.77</sup> )	0.7056 C <sub>5</sub> (sp <sup>1.78</sup> ) + 0.7056 C <sub>6</sub> (sp <sup>1.76</sup> )
N7-C14	0.6966 C <sub>5</sub> (sp <sup>1.85</sup> ) + 0.7175 C <sub>11</sub> (sp <sup>1.53</sup> )	0.6986 C <sub>5</sub> (sp <sup>1.79</sup> ) + 0.7156 C <sub>11</sub> (sp <sup>1.55</sup> )
N7-C14	0.7087 C <sub>5</sub> (p) + 0.7055 C <sub>11</sub> (p)	0.7090 C <sub>5</sub> (p) + 0.7052 C <sub>11</sub> (p)
N7-C14	0.7196 C <sub>17</sub> (sp <sup>1.62</sup> ) + 0.6943 C <sub>19</sub> (sp <sup>1.80</sup> )	0.7163 C <sub>17</sub> (sp <sup>1.64</sup> ) + 0.6978 C <sub>19</sub> (sp <sup>1.75</sup> )
N7-C14	0.7187 C <sub>17</sub> (p) + 0.6953 C <sub>19</sub> (p)	0.7195 C <sub>17</sub> (p) + 0.6945 C <sub>19</sub> (p)
N7-C14	0.7060 C <sub>11</sub> (sp <sup>1.97</sup> ) + 0.7082 C <sub>17</sub> (sp <sup>2.27</sup> )	0.7054 C <sub>11</sub> (sp <sup>1.95</sup> ) + 0.7088 C <sub>17</sub> (sp <sup>2.24</sup> )
N7-C14	0.7849 N <sub>7</sub> (sp <sup>1.80</sup> ) + 0.6196 C <sub>11</sub> (sp <sup>2.77</sup> )	0.7850 N <sub>7</sub> (sp <sup>1.77</sup> ) + 0.6195 C <sub>11</sub> (sp <sup>2.75</sup> )
N7-C14	0.7910 N <sub>7</sub> (sp <sup>1.88</sup> ) + 0.6118 C <sub>14</sub> (sp <sup>2.33</sup> )	0.7919 N <sub>7</sub> (sp <sup>1.86</sup> ) + 0.6107 C <sub>14</sub> (sp <sup>2.322</sup> )
N7-C14	0.6984 C <sub>14</sub> (sp <sup>1.86</sup> ) + 0.7157 C <sub>20</sub> (sp <sup>2.16</sup> )	0.6975 C <sub>14</sub> (sp <sup>1.85</sup> ) + 0.7166 C <sub>20</sub> (sp <sup>2.17</sup> )
N7-C14	0.7106 C <sub>17</sub> (sp <sup>2.21</sup> ) + 0.7036 C <sub>20</sub> (sp <sup>1.84</sup> )	0.7088 C <sub>17</sub> (sp <sup>2.20</sup> ) + 0.7054 C <sub>20</sub> (sp <sup>1.86</sup> )
N7-C14	0.8010 O <sub>10</sub> (sp <sup>1.51</sup> ) + 0.5987 C <sub>14</sub> (sp <sup>1.94</sup> )	0.8045 O <sub>10</sub> (sp <sup>1.44</sup> ) + 0.5939 C <sub>14</sub> (sp <sup>1.96</sup> )
N7-C14	0.8415 O <sub>10</sub> (p) + 0.5402 C <sub>14</sub> (p)	0.8376 O <sub>10</sub> (p) + 0.5463 C <sub>14</sub> (p)
N7-C14	0.8010 O <sub>10</sub> (sp <sup>1.99</sup> ) + 0.5987 C <sub>14</sub> (sp <sup>3.03</sup> )	0.8213 O <sub>12</sub> (sp <sup>1.97</sup> ) + 0.5705 C <sub>14</sub> (sp <sup>3.00</sup> )
N7-C14	0.5605 C <sub>4</sub> (sp <sup>3.54</sup> ) + 0.8281 O <sub>12</sub> (sp <sup>2.47</sup> )	0.5846 C <sub>4</sub> (sp <sup>3.42</sup> ) + 0.8254 O <sub>12</sub> (sp <sup>2.56</sup> )
N7-C14	0.6328 C <sub>20</sub> (sp <sup>2.07</sup> ) + 0.7743 N <sub>23</sub> (sp <sup>1.32</sup> )	0.6326 C <sub>20</sub> (sp <sup>2.00</sup> ) + 0.7745 N <sub>23</sub> (sp <sup>1.30</sup> )
N7-C14	0.6791 C <sub>20</sub> (p) + 0.7341 N <sub>23</sub> (p)	0.6778 C <sub>20</sub> (p) + 0.7353 N <sub>23</sub> (p)
N7-C14	0.7233 N <sub>22</sub> (sp <sup>2.27</sup> ) + 0.6906 N <sub>23</sub> (sp <sup>2.93</sup> )	0.7202 N <sub>22</sub> (sp <sup>2.25</sup> ) + 0.6938 N <sub>23</sub> (sp <sup>2.80</sup> )
N7-C14	0.6138 C <sub>18</sub> (sp <sup>2.45</sup> ) + 0.7723 N <sub>22</sub> (sp <sup>1.65</sup> )	0.6153 C <sub>18</sub> (sp <sup>2.41</sup> ) + 0.7883 N <sub>22</sub> (sp <sup>1.62</sup> )
N7-C14	0.6353 S <sub>8</sub> (sp <sup>4.38</sup> ) + 0.7723 C <sub>18</sub> (sp <sup>1.57</sup> )	0.6349 S <sub>8</sub> (sp <sup>4.02</sup> ) + 0.7651 C <sub>18</sub> (sp <sup>1.63</sup> )
N7-C14	0.8278 S <sub>5</sub> (p) + 0.5611 C <sub>18</sub> (p)	0.8291 S <sub>5</sub> (p) + 0.5591 C <sub>18</sub> (p)
N7-C14	0.6144 C <sub>18</sub> (sp <sup>2.10</sup> ) + 0.7723 N <sub>21</sub> (sp <sup>1.69</sup> )	0.6146 C <sub>18</sub> (sp <sup>2.05</sup> ) + 0.7888 N <sub>21</sub> (sp <sup>1.67</sup> )
N7-C14	0.6126 C <sub>15</sub> (sp <sup>2.70</sup> ) + 0.7904 N <sub>21</sub> (sp <sup>1.66</sup> )	0.6133 C <sub>15</sub> (sp <sup>2.68</sup> ) + 0.7899 N <sub>21</sub> (sp <sup>1.64</sup> )
N7-C14	1.96357 O <sub>12</sub> (sp <sup>1.63</sup> )	Unpaired electrons
N7-C14	1.84823 O <sub>12</sub> (p)	1.96232 O <sub>12</sub> (sp <sup>1.61</sup> )
N7-C14	1.98378 S <sub>8</sub> (sp <sup>0.22</sup> )	1.84485 O <sub>12</sub> (p)
N7-C14	1.86980 S <sub>8</sub> (sp <sup>99.99</sup> )	1.98517 S <sub>8</sub> (sp <sup>0.24</sup> )
N7-C14	1.97581 O <sub>10</sub> (sp <sup>0.66</sup> )	1.87217 S <sub>8</sub> (sp <sup>99.99</sup> )
N7-C14	1.85808 O <sub>10</sub> (sp <sup>99.99</sup> )	1.97502 O <sub>10</sub> (sp <sup>0.70</sup> )
N7-C14	1.66393 N <sub>7</sub> (p)	1.85401 O <sub>10</sub> (sp <sup>99.99</sup> )
N7-C14	1.92026 N <sub>23</sub> (sp <sup>2.15</sup> )	1.66484 N <sub>7</sub> (p)
N7-C14	1.58580 N <sub>22</sub> (p)	1.92038 N <sub>23</sub> (sp <sup>2.29</sup> )
N7-C14	1.63011 N <sub>21</sub> (p)	1.58989 N <sub>22</sub> (p)
N7-C14	1.99201 Cl <sub>37</sub> (sp <sup>0.21</sup> )	1.62631 N <sub>21</sub> (p)
N7-C14	1.97125 Cl <sub>37</sub> (p)	1.99224 Cl <sub>37</sub> (sp <sup>0.22</sup> )
N7-C14	1.93017 Cl <sub>37</sub> (p)	1.97100 Cl <sub>37</sub> (p)
N7-C14		1.93049 Cl <sub>37</sub> (p)



**Table 4** Calculated NBO analysis data for Zn(II) and Ni(II)-complexes calculated with 6-311G(d,p) basis sets.

Atom numbers	[Zn(HMICP) <sub>2</sub> ]	[Ni(HMICP) <sub>2</sub> ]
C3-C15	0.7007 C <sub>3</sub> (sp <sup>1.80</sup> ) + 0.7134 C <sub>15</sub> (sp <sup>1.84</sup> )	0.7004 C <sub>15</sub> (sp <sup>1.80</sup> ) + 0.7137 C <sub>16</sub> (sp <sup>1.82</sup> )
C15-C16	0.7177 C <sub>15</sub> (sp <sup>1.65</sup> ) + 0.6963 C <sub>16</sub> (sp <sup>1.94</sup> )	0.7153 C <sub>3</sub> (sp <sup>1.69</sup> ) + 0.6988 C <sub>15</sub> (sp <sup>1.88</sup> )
	0.7183 C <sub>15</sub> (p) + 0.6957 C <sub>16</sub> (p)	0.7070 C <sub>15</sub> (p) + 0.7072 C <sub>16</sub> (p)
C1-C3	0.7072 C <sub>1</sub> (sp <sup>1.78</sup> ) + 0.7070 C <sub>3</sub> (sp <sup>1.72</sup> )	0.7074 C <sub>1</sub> (sp <sup>1.78</sup> ) + 0.7068 C <sub>3</sub> (sp <sup>1.73</sup> )
	0.6981 C <sub>1</sub> (p) + 0.7160 C <sub>3</sub> (p)	0.6967 C <sub>1</sub> (p) + 0.7174 C <sub>3</sub> (p)
C9-C16	0.7095 C <sub>9</sub> (sp <sup>1.77</sup> ) + 0.7047 C <sub>16</sub> (sp <sup>1.82</sup> )	0.7083 C <sub>9</sub> (sp <sup>1.77</sup> ) + 0.7059 C <sub>16</sub> (sp <sup>1.77</sup> )
C1-C2	0.7069 C <sub>1</sub> (sp <sup>1.78</sup> ) + 0.7073 C <sub>2</sub> (sp <sup>1.58</sup> )	0.7066 C <sub>1</sub> (sp <sup>1.78</sup> ) + 0.7076 C <sub>2</sub> (sp <sup>1.58</sup> )
C2-C9	0.7095 C <sub>2</sub> (sp <sup>1.55</sup> ) + 0.7047 C <sub>9</sub> (sp <sup>1.79</sup> )	0.7086 C <sub>2</sub> (sp <sup>1.56</sup> ) + 0.7056 C <sub>9</sub> (sp <sup>1.78</sup> )
	0.7347 C <sub>2</sub> (p) + 0.67.84 C <sub>9</sub> (p)	0.7354 C <sub>2</sub> (p) + 0.6777 C <sub>9</sub> (p)
C2-Cl37	0.6881 C <sub>2</sub> (sp <sup>3.55</sup> ) + 0.7441 Cl <sub>37</sub> (sp <sup>4.38</sup> )	0.6990 C <sub>2</sub> (sp <sup>3.53</sup> ) + 0.7432 Cl <sub>37</sub> (sp <sup>4.40</sup> )
C6-C13	0.7072 C <sub>6</sub> (sp <sup>1.81</sup> ) + 0.7070 C <sub>13</sub> (sp <sup>1.61</sup> )	0.7078 C <sub>6</sub> (sp <sup>1.80</sup> ) + 0.7065 C <sub>13</sub> (sp <sup>1.61</sup> )
	0.7220 C <sub>6</sub> (p) + 0.6919 C <sub>13</sub> (p)	0.7210 C <sub>6</sub> (p) + 0.6930 C <sub>13</sub> (p)
C13-C19	0.7055 C <sub>9</sub> (sp <sup>1.75</sup> ) + 0.7087 C <sub>19</sub> (sp <sup>1.86</sup> )	0.7078 C <sub>9</sub> (sp <sup>1.74</sup> ) + 0.7064 C <sub>19</sub> (sp <sup>1.87</sup> )
C5-C6	0.7084 C <sub>5</sub> (sp <sup>1.78</sup> ) + 0.7058 C <sub>6</sub> (sp <sup>1.76</sup> )	0.7083 C <sub>5</sub> (sp <sup>1.79</sup> ) + 0.7059 C <sub>6</sub> (sp <sup>1.76</sup> )
C5-C11	0.6893 C <sub>5</sub> (sp <sup>1.79</sup> ) + 0.7158 C <sub>11</sub> (sp <sup>1.55</sup> )	0.6988 C <sub>5</sub> (sp <sup>1.78</sup> ) + 0.7153 C <sub>11</sub> (sp <sup>1.56</sup> )
	0.7147 C <sub>5</sub> (p) + 0.6994 C <sub>11</sub> (p)	0.7165 C <sub>5</sub> (p) + 0.6976 C <sub>11</sub> (p)
C17-C19	0.7162 C <sub>17</sub> (sp <sup>1.65</sup> ) + 0.6979 C <sub>19</sub> (sp <sup>1.78</sup> )	0.7175 C <sub>17</sub> (sp <sup>1.63</sup> ) + 0.6965 C <sub>19</sub> (sp <sup>1.80</sup> )
	0.7242 C <sub>17</sub> (p) + 0.6896 C <sub>19</sub> (p)	0.7310 C <sub>17</sub> (p) + 0.6824 C <sub>19</sub> (p)
C11-C17	0.7042 C <sub>11</sub> (sp <sup>1.97</sup> ) + 0.71.00 C <sub>17</sub> (sp <sup>2.26</sup> )	0.7050 C <sub>11</sub> (sp <sup>1.96</sup> ) + 0.7092 C <sub>17</sub> (sp <sup>2.27</sup> )
N7-C11	0.7833 N <sub>7</sub> (sp <sup>1.74</sup> ) + 0.6216 C <sub>11</sub> (sp <sup>2.71</sup> )	0.7837 N <sub>7</sub> (sp <sup>1.76</sup> ) + 0.6212 C <sub>11</sub> (sp <sup>2.71</sup> )
N7-C14	0.7940 N <sub>7</sub> (sp <sup>1.88</sup> ) + 0.6079 C <sub>14</sub> (sp <sup>2.29</sup> )	0.7933 N <sub>7</sub> (sp <sup>1.88</sup> ) + 0.6088 C <sub>14</sub> (sp <sup>2.27</sup> )
C14-C20	0.6946 C <sub>14</sub> (sp <sup>1.85</sup> ) + 0.7194 C <sub>20</sub> (sp <sup>2.18</sup> )	0.6928 C <sub>14</sub> (sp <sup>1.85</sup> ) + 0.7211 C <sub>20</sub> (sp <sup>2.41</sup> )
C17-C20	0.7103 C <sub>17</sub> (sp <sup>2.17</sup> ) + 0.7039 C <sub>20</sub> (sp <sup>1.85</sup> )	0.7099 C <sub>17</sub> (sp <sup>2.18</sup> ) + 0.7043 C <sub>20</sub> (sp <sup>1.68</sup> )
O10-C14	0.8056 O <sub>10</sub> (sp <sup>1.38</sup> ) + 0.5925 C <sub>14</sub> (sp <sup>1.91</sup> )	0.8046 O <sub>10</sub> (sp <sup>1.41</sup> ) + 0.5939 C <sub>14</sub> (sp <sup>1.93</sup> )
	0.8298 O <sub>10</sub> (p) + 0.5581 C <sub>14</sub> (p)	0.8319 O <sub>10</sub> (p) + 0.5549 C <sub>14</sub> (p)
O12-C13	0.8207 O <sub>12</sub> (sp <sup>1.98</sup> ) + 0.5713 C <sub>14</sub> (sp <sup>2.98</sup> )	0.8214 O <sub>12</sub> (sp <sup>1.98</sup> ) + 0.5703 C <sub>14</sub> (sp <sup>3.01</sup> )
C4-O12	0.5643 C <sub>4</sub> (sp <sup>3.42</sup> ) + 0.8255 O <sub>12</sub> (sp <sup>2.55</sup> )	0.5655 C <sub>4</sub> (sp <sup>3.40</sup> ) + 0.8247 O <sub>12</sub> (sp <sup>2.56</sup> )
C20-N23	0.6254 C <sub>20</sub> (sp <sup>1.99</sup> ) + 0.7803 N <sub>23</sub> (sp <sup>1.34</sup> )	0.6370 C <sub>20</sub> (sp <sup>2.02</sup> ) + 0.7794 N <sub>23</sub> (sp <sup>1.35</sup> )
	0.6363 C <sub>20</sub> (p) + 0.7714 N <sub>23</sub> (p)	0.6370 C <sub>20</sub> (p) + 0.7708 N <sub>23</sub> (p)
N22-N23	0.884 N <sub>22</sub> (sp <sup>2.85</sup> ) + 0.7253 N <sub>23</sub> (sp <sup>2.60</sup> )	0.6870 N <sub>22</sub> (sp <sup>3.02</sup> ) + 0.7266 N <sub>23</sub> (sp <sup>2.77</sup> )
C18-N22	0.6369 C <sub>18</sub> (sp <sup>1.81</sup> ) + 0.7709 N <sub>22</sub> (sp <sup>1.60</sup> )	0.6368 C <sub>18</sub> (sp <sup>1.85</sup> ) + 0.7711 N <sub>22</sub> (sp <sup>1.60</sup> )
	0.5880 C <sub>18</sub> (p) + 0.8089 N <sub>22</sub> (p)	
S8-C18	0.6994 S <sub>8</sub> (sp <sup>4.80</sup> ) + 0.7429 C <sub>18</sub> (sp <sup>2.04</sup> )	0.6638 S <sub>8</sub> (sp <sup>4.96</sup> ) + 0.7479 C <sub>18</sub> (sp <sup>2.02</sup> )
C18-N21	0.6125 C <sub>18</sub> (sp <sup>2.17</sup> ) + 0.7904 N <sub>21</sub> (sp <sup>1.61</sup> )	0.6132 C <sub>18</sub> (sp <sup>2.16</sup> ) + 0.7899 N <sub>21</sub> (sp <sup>1.62</sup> )
C15-N21	0.6125 C <sub>15</sub> (sp <sup>2.71</sup> ) + 0.7904 N <sub>21</sub> (sp <sup>1.63</sup> )	0.6154 C <sub>15</sub> (sp <sup>2.67</sup> ) + 0.7883 N <sub>21</sub> (sp <sup>1.65</sup> )
O12	1.96168 O <sub>12</sub> (sp <sup>1.61</sup> )	1.96204 O <sub>12</sub> (sp <sup>1.60</sup> )
	1.84425 O <sub>12</sub> (p)	1.84653 O <sub>12</sub> (p)
S8	1.97855 S <sub>8</sub> (sp <sup>0.35</sup> )	1.97355 S <sub>8</sub> (sp <sup>0.33</sup> )
	1.77965 S <sub>8</sub> (sp <sup>32.32</sup> )	1.75694 S <sub>8</sub> (sp <sup>94.52</sup> )
	1.75906 S <sub>8</sub> (sp <sup>15.72</sup> )	
O10	1.97489 O <sub>10</sub> (sp <sup>0.72</sup> )	1.97486 O <sub>10</sub> (sp <sup>0.71</sup> )
	1.84876 O <sub>10</sub> (P)	1.84603 O <sub>10</sub> (sp <sup>99.85</sup> )
N7	1.67306 N <sub>7</sub> (p)	1.67574 N <sub>7</sub> (sp <sup>99.99</sup> )
N23	1.86726 N <sub>23</sub> (sp <sup>2.38</sup> )	1.74252 N <sub>23</sub> (sp <sup>2.22</sup> )
N22	1.89774 N <sub>22</sub> (sp <sup>1.70</sup> )	1.90168 N <sub>22</sub> (sp <sup>1.71</sup> )
N21	1.62448 N <sub>21</sub> (p)	1.65347 N <sub>21</sub> (sp <sup>99.99</sup> )
Cl37	1.99230 Cl <sub>37</sub> (sp <sup>0.23</sup> )	1.99228 Cl <sub>37</sub> (sp <sup>0.23</sup> )
	1.97162 Cl <sub>37</sub> (p)	1.97125 Cl <sub>37</sub> (p)
	1.93191 Cl <sub>37</sub> (p)	1.93215 Cl <sub>37</sub> (p)

#### 4.1 UV Studies

H<sub>2</sub>MICP ligand was optimized at the basis set levels of B3LYP/6-31G(d,p), B3LYP/6-311G(d,p), B3LYP/6-311++G(d,p) and B3LYP/6-311++G(2d,2p). Excitation energies were obtained with time-dependent B3LYP (TDB3LYP TDHF) with the basis set of 6-31G(d,p), 6-311G(d,p), 6-311++G(d,p) and 6-311++G(2d,2p). Excitation energies of it is Zn(II) and Ni(II)-complexes optimized with B3LYP/6-311G(d,p) were obtained at the level of TDB3LYP/6-311G(d,p). The experimental and theoretical UV data of H<sub>2</sub>MICP ligand and its Zn(II) and Ni(II)-complexes are summarized in Table 5, while the excitation energies (eV) and oscillator strengths (f) are summarized in Table 6. Using calculation with the 6-311G(d,p), 6-311++G(d,p), 6-311++G(2d,2p) basis set, the peak which observed experimentally at 370 nm were obtained at 366, 367, 373 and 378 nm, respectively. This absorption is due to 1(HOMO) -1'(LUMO) and 4-1' electronic transition. HOMO(1) is composed of + 9.6% (S<sub>8</sub>) 3pz - 9.2% (C<sub>11</sub>) 2pz + 6.7% (C<sub>6</sub>) 2pz - 6.4% (O<sub>12</sub>) 2pz + 6.3% (C<sub>13</sub>) 2pz + 5.4% (S<sub>8</sub>) 4pz, while LUMO(1) is composed of + 16.6% (N<sub>23</sub>) 2pz + 8.4% (N<sub>23</sub>) 3pz - 7.4% (C<sub>20</sub>) 2pz - 6.0% (C<sub>20</sub>) 3pz + 5.4% (O<sub>10</sub>) 2pz - 5.1% (C<sub>14</sub>) 2pz, and HOMO-3(4) of + 10.7% (N<sub>22</sub>) 2pz - 9.7% (S<sub>8</sub>) 3pz + 9.3% (C<sub>19</sub>) 2pz - 6.9% (C<sub>20</sub>) 2pz + 6.4% (C<sub>19</sub>) 3pz + 5.4% (N<sub>22</sub>) 3pz atomic orbitals.

**Table 5** The experimental and theoretical UV data of H<sub>2</sub>MICP ligand and its Zn(II)-complex.

UV-visible spectrum data (nm)							
H <sub>2</sub> MICP							
Experimental	-	-	258	-	270	370	-
B3LYP/6-31G(d,p)	247	-	262	-	264	366	-
			260				
B3LYP/6-311G(d,p)	247	-	263	264	267	367	-
	241						
B3LYP/6-311++G(d,p)	245	-	265	267	270	373	-
				267			
B3LYP/6-311++G(2d,2p)	248	-	238	269	272	378	-
				267			
B3LYP/6-31G(d,p) (DMSO)	256	-	-	-	263	370	397
					261		
B3LYP/6-311G(d,p) (DMSO)	248	-	-	-	269	376	-
						327	
CEP	250	-	-	-	256	350	-
[Zn(HMICP) <sub>2</sub> ]							
Experimental	261	-	-	-	-	-	425
B3LYP/6-31G(d,p)	248	274	-	-	-	428	475
	245	265				409	470
	243	251				405	455
B3LYP/6-311G(d,p)	248	275	-	-	-	434	476
	246	256				409	473
		252				408	450
		251					



## 4.2 NMR Studies

B3LYP/6-31G(d,p), B3LYP/6-311G(d,p), B3LYP/6-311++G(d,p) and B3LYP/6-311++G(2d,2p) proton chemical shift calculations at both gas phase and DMSO solution were performed for H<sub>2</sub>MICP, while B3LYP/6-31G(d,p), and B3LYP/6-311G(d,p) proton chemical shift calculations at gas phase were conducted for [Zn(HMICP)<sub>2</sub>] (Table 7). The peak due to N22-H36 in the <sup>1</sup>H-NMR spectrum of H<sub>2</sub>MICP disappears in the spectrum of [Zn(HMICP)<sub>2</sub>], both in experimental measurement and theoretical calculation. The correlation between theoretical and experimental data was calculated as minimum 97% without chemical shifts of H31 and H35 which are N-H protons. The calculated chemical shifts have increased with basis set, and minimum values were observed with 6-31G(d,p), while maximum values were observed with 6-311++G(2d,2p) methods. The chemical shift values have increased in DMSO phase.

**Table 7** Experimental and theoretical proton chemical shifts for H<sub>2</sub>MICP, and [Zn(HMICP)<sub>2</sub>].

Atoms	Gas phase					DMSO		
	Exp. <sup>a</sup>	6-31G (d,p)	6-311G (d,p)	6-311++ G(d,p)	6-311++ G(2d,2p)	6-31G (d,p)	6-311G (d,p)	6-311++ G(2d,2p)
H <sub>2</sub> MICP*								
H34	7.35	7.20	7.39	7.51	7.59	7.38	7.58	7.59
H30	6.80	6.56	6.56	6.77	6.87	7.00	6.98	7.19
H29	6.90	6.58	6.70	6.66	6.81	7.18	7.32	7.14
H25	7.51	6.64	6.82	6.98	7.15	7.22	7.70	7.50
H33	7.51	9.79	9.90	9.95	10.31	9.62	7.74	8.76
H24	7.45	7.19	7.34	7.50	7.60	7.67	7.88	7.75
H32	7.45	7.28	7.44	7.52	7.61	7.64	7.86	7.79
H35	12.82	9.38	9.41	9.70	10.00	9.53	9.02	9.96
H36	11.05	12.66	12.44	12.63	13.18	12.98	12.90	13.24
H31	10.76	6.06	6.07	6.36	6.66	8.04	8.10	7.32
H26	3.75	3.58	3.59	3.66	3.74	3.70	3.66	3.84
H27	3.75	4.01	4.08	4.14	4.09	4.04	4.07	4.11
H28	3.75	3.58	3.59	3.66	3.74	3.69	3.67	3.84
Regr.	-	0.97	0.97	0.97	0.97	0.98	0.99	0.98
[Zn(HMICP) <sub>2</sub> ]								
H34	6.85-7.66	7.14	6.88	-	-	-	-	-
H30	-	6.37	6.14	-	-	-	-	-
H29	-	6.38	6.27	-	-	-	-	-
H25	-	6.50	6.45	-	-	-	-	-
H33	-	10.20	10.03	-	-	-	-	-
H24	-	7.11	7.01	-	-	-	-	-
H32	-	7.50	7.42	-	-	-	-	-
H35	10,81	6.87	6.92	-	-	-	-	-
H31	10.65	6.03	5.86	-	-	-	-	-
H26	3.70	3.62	3.46	-	-	-	-	-
H27	3.62	3.28	3.04	-	-	-	-	-
H28	3.30	3.25	3.00	-	-	-	-	-

\*without H31 and H35 for H<sub>2</sub>MICP, a: in [14]

**Table 8** Experimental and theoretical vibrational assignments of H<sub>2</sub>MICP carried out with B3LYP method and 6-31G(d,p), 6-311G(d,p), 6-311++G(d,p), and 6-311++G(2d,2p) basis sets.

Exp.	6-31G(d,p)		6-311G(d,p)		6-311++G (d,p)		6-311++G (2d,2p)		Assignments
	Freq.	Int. <sup>1</sup>	Freq.	Int.	Freq.	Int.	Freq.	Int.	
3284	3663	75	3644	75	3642	78	3648	77	v(N <sub>7</sub> H) <sub>indole</sub>
3254	3513	100	3500	108	3504	104	3517	98	v(N <sub>21</sub> H) <sub>thio</sub>
3230	3426	104	3422	98	3421	96	3431	100	v(N <sub>22</sub> H) <sub>thio</sub>
-	3252	16	3236	14	3232	16	3238	18	v(C <sub>15</sub> -H) <sub>ringC</sub> , v(C <sub>9</sub> H) <sub>ringC</sub>
-	3200	12	3182	12	3181	10	3187	10	v(C <sub>6</sub> -H) <sub>ringA</sub> , v(C <sub>3</sub> H) <sub>ringA</sub>
-	3178	10	3162	9	3161	9	3168	8	v(C <sub>1</sub> -H) <sub>ringC</sub> , v(C <sub>3</sub> H) <sub>ringC</sub>
-	3153	22	3136	21	3137	19	3142	17	v(CH <sub>3</sub> ) <sub>met</sub>
-	3078	41	3058	42	3062	37	3071	33	v(CH <sub>3</sub> ) <sub>met</sub>
-	3016	69	3000	68	3004	69	3015	64	v(CH <sub>3</sub> ) <sub>met</sub>
1697	1789	252	1773	284	1758	325	1747	307	v(C-O), δ(N <sub>22</sub> H)
1621	1690	8	1678	10	1674	9	1671	8	v(CC) <sub>ringA com.</sub> <sup>2</sup> , δ(N <sub>7</sub> H), δ(OCH <sub>3</sub> ) <sub>met</sub> , v(N <sub>20</sub> C <sub>23</sub> )
-	1654	9	1640	4	1636	4	-	-	v(CC) <sub>ringC com.</sub> , δ(N <sub>14</sub> H), v(N <sub>20</sub> C <sub>23</sub> )
1595	1649	98	1636	112	1633	99	1636	14	v(CC) <sub>ringA com.</sub> , δ(N <sub>8</sub> H), v(OCH <sub>3</sub> ) <sub>met</sub> , v(N <sub>20</sub> C <sub>23</sub> )
1573	1644	133	1635	199	1633	174	1632	188	v(CC) <sub>ringC com.</sub> , δ(N <sub>21</sub> H)
1541	1641	232	1626	196	1620	218	1630	65	v(CC) <sub>com.</sub> , δ(N <sub>7</sub> H), v(N <sub>20</sub> C <sub>23</sub> ), δ(N <sub>22</sub> H)
-	-	-	-	-	-	-	1610	213	v(CC) <sub>ringA com.</sub> , δ(N <sub>7</sub> H), v(N <sub>20</sub> C <sub>23</sub> )
1487	1590	771	1582	675	1578	680	1577	634	v(CC) <sub>ringC com.</sub> , δ(N <sub>21</sub> H), v(N <sub>21</sub> C <sub>18</sub> )
1481	1536	154	1527	186	1524	172	1529	178	v(CC) <sub>ringC com.</sub> , δ(N <sub>21</sub> H), δ(N <sub>22</sub> H)
1430	1532	97	1521	218	1518	213	1517	38	v(CC) <sub>ringC com.</sub> , δ(N <sub>21</sub> H), δ(N <sub>22</sub> H)
1397	1527	240	1518	166	1514	157	1516	221	v(CC) <sub>ringA</sub> , δ(N <sub>22</sub> H), s-cis-methoxy, v(C <sub>18</sub> S <sub>8</sub> )
1383	1516	270	1504	215	1502	246	1508	304	v(CC) <sub>ringA</sub> , δ(N <sub>22</sub> H), s-cis-methoxy
-	1505	6	1493	8	1493	10	1500	9	s-cis-methoxy
1308	1500	31	1489	30	1487	31	1488	30	v-isatin, δ(N <sub>22</sub> H), s-cis (CH <sub>3</sub> ) <sub>met</sub>
1293	1485	91	1475	77	1472	63	1476	67	v(CC) <sub>ringA com.</sub> , s-cis-methoxy
-	1446	13	1437	7	1433	6	1436	7	v(CC) <sub>ringC com.</sub> , δ(N <sub>21</sub> H), δ(N <sub>22</sub> H)
1273	1436	32	1427	34	1424	32	1420	35	v(CC) <sub>ringC com.</sub> , δ(N <sub>7</sub> H)
1240	1405	323	1394	383	1391	385	1388	414	δ(N <sub>21</sub> C <sub>18</sub> ), v(N <sub>22</sub> C <sub>18</sub> N <sub>21</sub> )
-	-	-	1339	7	1338	8	1341	10	v(CC) <sub>ringC com.</sub>
1195	1339	176	1327	164	1325	148	1327	151	v(CC) <sub>ringC com.</sub> , δ(N <sub>7</sub> H), s-cis-OCH <sub>3</sub>
-	1330	3	1324	5	1322	8	1320	28	v(CC) <sub>ringC com.</sub>
1155	1324	83	1311	61	1309	57	1310	35	Δ-isatin
1112	1306	77	1293	73	1290	78	1289	65	v-isatin, δ(N <sub>22</sub> H), s-cis-OCH <sub>3</sub>

<sup>1</sup>Intensity, <sup>2</sup>combination

### 4.3 IR Studies

Experimental and theoretical vibrational assignments of H<sub>2</sub>MICP were carried out with the aid of B3LYP method and 6-31G(d,p), 6-311G(d,p), 6-311++G(d,p), and 6-311++G(2d,2p) basis sets as shown in Table 8. For studied basis sets, correlation coefficients were found as 0.971, and 0.969. Absorption bands at 3650, 3294 and 3192 cm<sup>-1</sup> were identified. B3LYP results showed that the vibrational modes of  $\nu(\text{N}_7\text{H})_{\text{indole}}$ ,  $\nu(\text{N}_{22}\text{H})_{\text{thio}}$ , and  $\nu(\text{N}_{21}\text{H})_{\text{thio}}$ , are as follow: 3663, 3513, and 3426 cm<sup>-1</sup> for 6-31G(d,p); 3644, 3500, and 3422 cm<sup>-1</sup> for 6-311G(d,p); 3642, 3504, and 3421 cm<sup>-1</sup> for 6-311++G(d,p); 3648, 3517, and 3431 cm<sup>-1</sup> for 6-311++G(2d,2p). The band at 3294 cm<sup>-1</sup> belonging to  $\nu(\text{N}_{22}\text{H})$  disappears in its zinc(II) and nickel(II)-complexes. The other two bands nearly remained unchanged in both zinc(II) and nickel(II)-complexes. According to theoretical result, the absorption between 3252 and 2936 cm<sup>-1</sup> can be assigned to the vibrational modes of  $\nu(\text{C-H})_{\text{ringC}}$ ,  $\nu(\text{C-H})_{\text{ringA}}$ ,  $\nu(\text{CH}_3)_{\text{met}}$ . In the infrared spectrum of C=O for H<sub>2</sub>MICP, we observe the band at 1697 cm<sup>-1</sup>. By means of the DFT procedure with B3LYP/6-31G(d,p) (1789 cm<sup>-1</sup>), B3LYP/6-311G(d,p) (1773 cm<sup>-1</sup>), B3LYP/6-311++G(d,p) (1758 cm<sup>-1</sup>), and B3LYP/6-31G(2d,2p) (1747 cm<sup>-1</sup>) basis sets, we can assign the band experimentally observed at 1697 cm<sup>-1</sup> (IR) is the  $\nu(\text{C=O})$  vibrational mode. The band observed at 1697 cm<sup>-1</sup> for H<sub>2</sub>MICP appeared at 1694 cm<sup>-1</sup> for its zinc(II)-complex, indicating that the C=O group is not involved in coordination and at 1670 cm<sup>-1</sup> for its nickel(II)-complexes indicating that C=O group is involved in coordination. The absorption at about 850 cm<sup>-1</sup> for H<sub>2</sub>MICP is assignable to the vibrational modes involving the C=S group. This absorption is assigned at about 820 cm<sup>-1</sup> for its Zn(II)-complex and 818 cm<sup>-1</sup> for its Ni(II)-complex due to transfer of charge from sulfur atom to the metal.

## 5 Conclusions

New Zn(II) and Ni(II)-complexes of H<sub>2</sub>MICP have been synthesized, and its theoretical study has also been conducted. For the HOMO, it was found that the main contributions due to the thiosemicarbazone group and isatin group. Vibrational study using B3LYP calculations showed that the disappearance of the  $\nu(\text{N}_{22}\text{H})$  band of H<sub>2</sub>MICP ligand indicates the deprotonation of the group in coordination. Analysis of experimental and theoretical UV, IR and NMR data of H<sub>2</sub>MICP ligand and its Zn(II) and Ni(II)-complexes showed that theoretical calculations are in line to supporting the experimental results.

## Acknowledgement

Authors thank TUBITAK ULAKBIM, High Performance and Grid Computing Center (TR-Grid e-Infrastructure) for performing the numerical calculations.

## References

- [1] Pandeya, S.N., Sriram, D., Nath, G. & De Clercq, E., *Synthesis, Antibacterial, Antifungal and Anti HIV activities of Schiff and Mannich bases derived from isatin derivatives and N-[4-(4-chlorophenyl) hiazol-2-yl] thiosemicarbazide*, Eur. J. Pharma. Sci., **9**, pp. 25-31, 1999.
- [2] Bauer, D.J. & Sadler P.W., *The structure-activity relationships of the antiviral chemotherapeutic activity of isatin  $\beta$ -thiosemicarbazone*, Brit. J. Pharmacol. **15**, pp. 101-110, 1960.
- [3] Levinson, W. & Helling, R., *Inactivation of lambda phage infectivity and lambda deoxyribonucleic acid transfection by N-methyl-isatin  $\beta$ -thiosemicarbazone-copper complexes*, Antimicrob. Agents Chemother., **9**, pp. 160-163, 1976.
- [4] Qu, J.Q., Sun, G.C., Wang, L.F. & Qu, L., *Synthesis, Characterization, And Biological Activities of Some Transition metal(II) Complexes with the Thiosemicarbazone derived from 4-[1-(4-methylphenylsulfonyl)-1 H -indol-3-yl]but-3-en-2-one*, Chem. Mat. Sci., **60**(3), pp. 214-219, 2006.
- [5] Karali, N., Gürsoy, A., Kandemirli, F., Shvets, N., Kaynak, .F.B., Özbey, S., Kovalishyn, V. & Dimoglo, A., *Synthesis and Structure-Antituberculosis Activity Relationship of 1H-indole-2,3-dione derivatives*, Bioorg. Med. Chem., **15**(17), pp. 5888-5904, 2007.
- [6] Saracoglu, M., Kandemirli, F., Kovalishyn, V., Arslan, T. & Ebenso, E.E., *ETM-ANN Approach Application for Thiobenzamide and Quinolizidine Derivatives*, J. Biomed. and Biotech., **2010**, doi:10.1155/2010/693031, 12 pages, 2010.
- [7] Kandemirli, F., Arslan, T., Koksoy, B. & Yilmaz, M., *Synthesis, characterization and theoretical calculations of 5-methoxyisatin-3-thiosemicarbazone derivatives*, J. Chem. Soc. Pak., **31**(3), pp. 498-504, 2009.
- [8] Parr, R.G. & Yang W., *Density Functional Theory Of Atoms And Molecules*, Oxford University Press. Oxford. 1989.
- [9] A.D. Becke, *A New Mixing of Hartree-Fock and Local Density-Functional Theories*, J. Chem. Phys., **98**, pp. 1372-1377, 1993.
- [10] Frisch, M.J., Trucks, G.W., Schlegel, H.B., Scuseria, G.E., Robb, M.A., Cheeseman, J.R., Scalmani, G., Barone, V., Mennucci, B., Petersson, G.A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H.P., Izmaylov, A.F., Bloino, J., Zheng, G., Sonnenberg, J.L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery, Jr. J.A., Peralta, J.E., Ogliaro, F., Bearpark, M., Heyd, J.J., Brothers, E., Kudin, K.N., Staroverov, V.N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J.C., Iyengar, S.S., Tomasi, J., Cossi, M., Rega, N., Millam, J.M., Klene, M., Knox, J.E., Cross, J.B., Bakken, V., Adamo, C.,

- Jaramillo, J., Gomperts, R., Stratmann, R.E., Yazyev, O., Austin, A.J., Cammi, R., Pomelli, C., Ochterski, J.W., Martin, R.L., Morokuma, K., Zakrzewski, V.G., Voth, G.A., Salvador, P., Dannenberg, J.J., Dapprich, S., Daniels, A.D., Farkas, Ö., Foresman, J.B., Ortiz, J.V., Cioslowski, J. & Fox, D.J., *Gaussian 09, Revision A.1*, Gaussian, Inc., Wallingford CT, 2009.
- [11] Kandemirli, F., Arslan, T., Karadayı, N., Ebenso, E.E. & Koksoy, B., *Synthesis and Theoretical Study of 5-methoxyisatin-3-(N-cyclohexyl)-Thiosemicarbazone and Its Ni(II) and Zn(II) complexes*, J. Mol. Struc., **938**, pp. 89–96, 2009.
- [12] Gorelsky, S.I., *AOMix: Program for Molecular Orbital Analysis*, University of Ottawa, Ottawa, Canada, 2009, <http://www.sg-chem.net>.
- [13] Gorelsky, S.I. & Lever, A.B.P., *Electronic Structure And Spectra Of Ruthenium Diimine Complexes By Density Functional Theory and INDO/S. Comparison of the Two Methods*, J. Organomet. Chem., **635**, pp. 187-196, 2001.
- [14] Weinhold, F., *Natural Bond Orbital Analysis Programs Theoretical Chemistry Institute and Department of Chemistry*, University of Wisconsin, Madison, Wisconsin, 53706.